



RF-3615

M. Sc. (Part - II) Examination  
April / May - 2010  
Advance Integral Transforms - 5006

Time : Hours]

[Total Marks :

Instructions :

(1)

नीचे दशांशवले निशानीवाणी विगतो उत्तरवळी पर अवश्य कभवली.  
Fillup strictly the details of signs on your answer book.

Name of the Examination :  
M. SC. - 2

Name of the Subject :  
ADVANCE INTEGRAL TRANSFORMS - 5006

Subject Code No. : 3 6 1 5 Section No. (1, 2,.....) : NIL

Seat No. :

Student's Signature

- (2) All questions are compulsory.  
 (3) Notations used are standard.  
 (4) Figure at the right end of the first line of each question indicate full marks.

Q.1	(a)	State and prove inverse theorem for Mellin transform.	[6]
	(b)	Define Inverse Mellin transform. State and prove convolution theorem for Mellin transform.	[4]
	(c)	Prove that $M[f^n(x):s] = \frac{(-1)^n(s-1)!}{[(s-n)-1]!} f^*(s-n)$ .	[4]
OR			
Q.1	(a)	Find the potential $\phi(r, \theta)$ that satisfies the linear equation $r^2\phi_{rr} + r\phi_r + \phi_{\theta\theta} = 0$ in an infinite wedge $0 < r < \infty; -\alpha < \theta < \alpha$	[6]
	(b)	with boundary conditions $\phi(r, \alpha) = f(r); \phi(r, -\alpha) = g(r); 0 \leq r < \infty, \phi(r, \theta) \rightarrow 0$ as $r \rightarrow \infty \forall \theta \in (-\alpha, \alpha)$ .	[4]
	(c)	In usual notation prove that $M\left[\frac{\overline{a}}{(1+x)^a}:s\right] = \overline{a-s} \overline{s}$ and deduce $M\left[\frac{1}{(1+x)}:s\right] = \pi \operatorname{cosec}(\pi s)$ .	[4]
Q.2	(a)	Let $\overline{f}_n(k)$ and $\overline{f}'_n(k)$ be the Hankel transform of order n of functions f(r) and f'(r) respectively then $\overline{f}'_n(k) = -\frac{k}{2n}[(n+1)\overline{f}_{n-1}(k) - (n-1)\overline{f}_{n+1}(k)]$ .	[6]
	(b)	Find Hankel transform of order n of $\frac{d^2 f}{dr^2} + \frac{1}{r} \frac{df}{dr} - \frac{n^2}{r^2} f$ .	[4]
	(c)	State and prove Parseval's relation for Hankel transform.	[4]

OR

Q.2	(a)	The magnetic potential $v$ of the circular disc of radius 'a' and strength ' $\omega$ ' magnetized parallel to axis satisfies Laplace equation and it is equal to $2\pi\omega$ on the disc itself and it vanishes at the exterior points in the plane of a disc. Show that at point (r,z) and $z > 0$ $v = 2\pi\omega a \int_0^\infty e^{-kz} J_0(kr) J_1(ak) dk$ .	[6]
	(b)	Find Hankel transform of $f(r) = \begin{cases} a^2 - r^2; 0 < r \leq a \\ 0; r \geq a \end{cases}; n = 0$ .	[4]
	(c)	State and prove Change of scale property for Hankel transform.	[4]
Q.3	(a)	Use Finite Hankel transform to solve differential equation $\frac{\partial^2 v}{\partial r^2} + \frac{1}{r} \frac{\partial v}{\partial r} = \frac{1}{k_1} \frac{\partial v}{\partial t}; 0 \leq r \leq 1; t > 0$ where $v = v_0 = \text{constant for } r = 1$ $v = 0$ for $t = 0$ and $0 \leq r < 1$ .	[6]
	(b)	Show that Finite Hankel transform, $H_0^*(r^2) = \frac{a^3}{k} J_1(ka) - \frac{2a^2}{k^2} J_2(ka)$ .	[4]
	(c)	Find the Finite Hankel transform of $r^{n-1}$ where $r J_{n-1}(kr)$ is the kernel of transform ( $n > 1$ ).	[4]
<b>OR</b>			
Q.3	(a)	Show that $\int_0^1 \frac{J_n(ar)}{J_n(a)} r J_n(kr) dr = \frac{k}{a^2 - k^2} J'_n(k)$ where $J_n(k) = 0$ .	[6]
	(b)	Prove that Finite Hankel transform, $H_n^* \left[ \frac{d^2 f}{dr^2} + \frac{1}{r} \frac{df}{dr} \right] = \frac{k}{2} \left[ H_{n+1}^* \left( \frac{df}{dr} \right) - H_{n-1}^* \left( \frac{df}{dr} \right) \right]$	[4]
	(c)	Find the Finite Hankel transform of order zero of a constant function.	[4]
Q.4	(a)	Solve the Initial value problem $u_{n+2} - 5u_{n+1} + 6u_n = 2^n; u_0 = 1$ and $u_1 = 0$ using Z-transform.	[6]
	(b)	If $Z[u_n] = \frac{2z^2 + 3z + 4}{(z-1)^3}$ . Find $u_1$ and $u_2$ .	[4]
	(c)	Find (i) $Z^{-1} \left[ \frac{3z^2 - z}{(z-1)(z-2)^2} \right]$ and (ii) $Z^{-1} \left[ \frac{z}{(z-1)(z-2)} \right]$ .	[4]
<b>OR</b>			
Q.4	(a)	Prove that $Z[n^p] = -z \frac{d}{dz} [Z(n^{p-1})]$ where $p > 0$ is any integer and hence deduce $Z[n] = \frac{z}{(z-1)^2}$ and $Z[n^2] = \frac{z^2 + z}{(z-1)^3}$ .	[6]
	(b)	State and prove Shifting theorem for Z-transform.	[4]
	(c)	Find $Z[\cos(n\theta)]$ .	[4]
Q.5	(a)	Define Hilbert Transform. Also derive the formula for Inverse Hilbert transform.	[6]
	(b)	Find the Hilbert transform of $f(t) = \cos(\omega t)$ .	[4]
	(c)	If $f(t) = e^{-at}$ find Generalized Stieltjes transform of $f(t)$ .	[4]

**OR**

Q.5	(a)	Find the solution of the integral equation given by $\lambda \int_0^\infty \frac{f(t)}{t+x} dt = f(x)$ where $\lambda$ is a real parameter, using Stieltjes transform.	[6]
	(b)	Find Hilbert transform of $f(t) = \frac{t}{t^2 + a^2}; a > 0$ .	[4]
	(c)	Find Stieltjes transform of (i) $f(t) = (t+a)^{-1}$ and (ii) $f(t) = t^{\alpha-1}$ .	[4]